INDEX TO VOLUME 72 (1988)[1]
AUTHORS[2]

Acevedo-Alfaro, J.
Bacteria associated with the West Indian fruit fly *Anastrepha obliqua* (Macquart) (Diptera-Tephritidae), 72(3), 489 (RN).

Almodóvar, L.
Período crítico de competencia de las malezas en el tomate de trasplante, 72(2), 291.

Almodóvar-Vega, L.
*Evaluation of diquat for postemergence weed control in pumpkin (*Curcubita moschata*), 72(2), 285.

*Allelopathic effects of seven weed species to pumpkin (*Curcubita moschata*) under greenhouse conditions, 72(3), 495 (RN).

*Eggplant (*Solanum melongena* L.) transplants and exudates from roots of seven weeds, 72(3), 495 (RN).

Armstrong, A. M.
*Chemical control of *Liriomyza sativae* and of Lepidoptera larvae on tomato, 72(1), 169 (RN).

†Light trap studies of *Heliothis virescens* (Fabricius) and *Etiella zinckenella* (Treitschke) on pigeon pea (*Cajanus cajan* Millsp), 72(4), 557.

Badillo-Feliciano, J.
Yield response of vegetables to tillage techniques on a vertisol of the semiarid region of Puerto Rico, 72(1), 119.

Barbosa, P.
Relative susceptibility of *Crotalaria* spp. to attack by *Etiella zinckenella* in Puerto Rico, 72(1), 147.

Notes on the natural enemies of *Etiella zinckenella* in Puerto Rico, 72(1), 153.

The sorghum midge, *Contarinia sorghicola*, a sorghum pest newly introduced to Puerto Rico, 72(1), 161 (RN).

Beauchamp de Caloni, I.
*Freezing and shelf life of sweet potato and citron bars, 72(1), 171 (RN).

Sensory evaluation of cooked new sweet potato selections, 72(2), 265.

Evaluación de variedades de uva de mesa (*Vitis vinifera* L.) cosechadas en la Subestación de Fortuna en 1986, 72(2), 335 (RN).

The Lacknau plantain: A high yielding cultivar with field resistance to the corm weevil, *Cosmopolites sordidus* (Germar), 72(3), 353.

*Senior author.
†Sole author.
RN Research note.

[1]An author's name may appear in more than one form. Although all first names are abbreviated, an author may appear with one or two initials. Last names may appear single or compound. When compound, they are hyphenated regardless of how they appear in the original.

Alphabetically, names with a single initial precede two-initial ones (Y. Badillo precedes Y. A. Badillo); single names precede hyphenated ones (Abreu, Z. precedes Abreu-Abadía, A.).

Articles under one name are organized chronologically (those in volume I precede those in volume II; those in number 1 precede those in number 2 and those on page 1 precede those on page 2).

Beaver, J. S.
Heritabilities and phenotypic correlations for seed yield and seed yield components of beans, 72(2), 301.
*Dry beans and varying cultural practices at two locations on the coastal plain of Puerto Rico, 72(4), 521.
Beinroth, F. H.
Yield response of vegetables to tillage techniques on a Vertisol of the semiarid region of Puerto Rico, 72(1), 119.
Bellerive, P.
Cost of a drip irrigation system for vegetable farming in Puerto Rico, 72(1), 31.
Bendezú, P.
*Watercress cultivation sites and their relationship to fascioliasis in Puerto Rico, 72(3), 405.
Biascoechea, M. L.
Tanier (*Xanthosoma* spp.) propagation in vitro, 72(3), 413.
Bonilla, C.
Watercress cultivation sites and their relationship to fascioliasis in Puerto Rico, 72(3), 405.
Bosques, A.
Salt and sugar levels for effective plant regeneration from in vitro cultured tissues of cocoyam, 72(4), 515.
Cacho, E
*Pilot plant fermentation of high test molasses compared with blackstrap molasses using factorial design, 72(1), 9.
Evaluation of a commercial yeast nutrient for high test molasses fermentation, 72(1), 173 (RN).
*The use of distillers' dry yeast as inoculum for rum fermentation on pilot plant scale, 72(3), 445.
*Pilot plant scale evaluation of a commercial yeast nutrient for high test molasses fermentation, 72(4), 509.
Caloni (de), 1. B.
Shelf life and acceptability of hot water-treated mangoes, 72(3), 469.
Cardona-Castro, C.
Agronomic performance of hard endosperm opaque-2 corn populations in Puerto Rico, 72(2), 309.
Carpintero, J. D.
*Contributions to the knowledge of American Ectrichodiinae. I. Notes about *Rhiginia* (Hemiptera:Reduviidae), 72(2), 251.
Cedeño-Maldonado, A.
*Effect of dwarfing rootstocks on tree size and yield of selected mango varieties, 72(1), 1.
Dwarfing effect of interstems and growth and yield components of mango, 72(4), 501.
*Salt and sugar levels for effective plant regeneration from in vitro cultured tissues of cocoyam, 72(4), 515.
Chao de Báez, C.
Temperature versus elevation relationships for Puerto Rico, 72(3), 449.
Nutrient uptake and growth characteristics of nitrogen fertigated sweet peppers under drip irrigation and plastic mulch, 72(4), 575.
Tres niveles de riego y crecimiento y rendimiento del maíz dulce (*Zea mays* v. Suresweet) regado por goteo, 72(4), 565.
Colberg-Rivera, 0.
†Seed production of asparagus (*Asparagus officianalis*) in Puerto Rico, 72(4), 619 (RN).
Corchado-Juarbe, N.
Storage and digestibility, voluntary intake and chemical components of hay of five tropical grasses, 72(4), 531.
Cortés-Monllor, A.
†Bacterial leaf spot of coffee, 72(4), 621 (RN).
Coscarón, M. C.
*Contribution to the knowledge of *Rasahus* Amyot and Serville, 1843 (Hemiptera:Reduviidae): Description of a new species, 72(2), 247.
Cotte, 0.
*Some host plants of leafhopper, *Empoasca* spp. (Homoptera:Cicadellidae), found on pigeon pea (*Cajanus cajan* L.) fields, 72(3), 499 (RN).
Crespo-Ruiz, M.
Root distribution of nitrogen fertigated sweet peppers under drip irrigation, 72(1), 51.
*Nutrient uptake and growth characteristics of nitrogen fertigated sweet peppers under drip irrigation and plastic mulch, 72(4), 575.
Cruz, C.
Chemical control of *Liriomyza sativae* and of Lepidoptera larvae on tomato, 72(1), 169 (RN).

*Senior author.
†Sole author.
RN Research note.

Some host plants of leafhopper, *Empoasca* spp. (Homoptera:Cicadellidae), found in pigeon pea (*Cajanus cajan* L.) fields, 72(3), 499 (RN).

Díaz, N.
*Shelf life and acceptability of hot water-treated mangoes, 72(3), 469.

Díaz-Rivera, M.
Evaluation of diquat for postemergence weed control in pumpkin (*Curcubita moschata*), 72(2), 285.

Escudero, J.
Bacteria associated with the West Indian fruit fly *Anastrepha obliqua* (Macquart) (Diptera:Tephritidae), 72(3), 489 (RN).

Fernández-Coll, F.
*Refrigeration before incubation and its effect on recovery of two types of bacteria from food, 72(4), 617 (RN).

Fernández-VanCleve, J.
Yield and nutritive value of hay from five tropical grasses at three harvesting intervals, 72(1), 109.

Confinement feeding of dairy cows based on stargrass as green chopped fodder or hay, 72(2), 231.

Fierro, C.
Salt and sugar levels for effective plant regeneration from in vitro cultured tissues of cocoyam, 72(4), 515.

Figueroa, L. A.
Yam protection. II. Antracnose, yield, and profit of monocultures and interplanting, 72(2), 179.

Fontanet, E.
Pilot plant fermentation of high test molasses compared with blackstrap molasses using factorial design, 72(1), 9.

Yield and nutritive value of hay from five tropical grasses at three harvesting intervals, 72(1), 109.

Pilot plant scale evaluation of a commercial yeast nutrient for high test molasses fermentation, 72(4), 509.

Frame, A. D.
Watercress cultivation sites and their relationship to fascioliasis in Puerto Rico, 72(3), 405.

Frame, E. L.
Watercress cultivation sites and their relationship to fascioliasis in Puerto Rico, 72(3), 405.

*Senior author.
†Sole author.
RN Research note.

Franqui, R. A.
*Host plants of pentatomids affecting rice fields in Puerto Rico, 72(3), 365.

*Natural enemies of pentatomids affecting rice fields in Puerto Rico, 72(3), 371.

Gallardo-Covas, F.
†Distribution of the coffee leaf miner, *Leucoptera coffeella*, and its parasitoids in the canopy of coffee, *Coffea arabica*, in Puerto Rico, 72(1), 141.

*Susceptibilidad de 53 cultivares de café, *Coffea arabica*, al daño del minador del café, *Leucoptera coffeella*, en Puerto Rico, 72(1), 165 (RN).

†Faunal survey of the coffee leaf miner, *Leucoptera coffeella*, parasitoids in Puerto Rico, 72(2), 255.

González, E. A.
Temperature versus elevation relationships for Puerto Rico, 72(3), 449.

Requisitos de riego para plátano en siete regiones ecológicas de Puerto Rico, 72(4), 599.

Requisitos de riego para sorgo en las costas sur y norte de Puerto Rico, 72(4), 585.

Seasonal consumptive use of water by Bell and Cubanelle peppers in semiarid andhumid coastal sites in Puerto Rico, 72(4), 609.

González-Figueroa, W.
Bacteria associated with the West Indian fruit fly *Anastrepha obliqua* (Macquart) (Diptera:Tephritidae), 72(3), 489 (RN).

González-Fuentes, E. A.
*Estimación de los requisitos de riego para maíz dulce (*Zea mays* cv. Suresweet) en la costa sur de Puerto Rico, 72(2), 277.

*Tres niveles de riego, crecimiento y rendimiento del maíz dulce (*Zea mays* cv. Suresweet) regado por goteo, 72(4), 565.

González-Muñiz, W.
Susceptibilidad de 53 cultivares de café, *Coffea arabica*, al daño del minador del café, *Leucoptera coffeella*, en Puerto Rico, 72(1), 165 (RN).

Goyal, M. R.
Cost of a drip irrigation system for vegetable farming in Puerto Rico, 72(1), 31.

*Labor-input requirements for experimental production of drip irrigated vegetables, 72(1), 41.

*Root distribution of nitrogen fertigated sweet peppers under drip irrigation, 72(1), 51.

†Potential evapotranspiration for the south coast of Puerto Rico with the Hargreaves-Samani technique, 72(1), 57.

†Potential evapotranspiration for Vieques Island, Puerto Rico, with the Hargreaves-Samani model, 72(1), 177 (RN).

Estimación de los requisitos de riego para maíz dulce (*Zea mays* cv. Suresweet) en la costa sur de Puerto Rico, 72(2), 277.

*Temperature versus elevation relationships for Puerto Rico, 72(3), 449.

Nutrient uptake and growth characteristics of nitrogen fertigated sweet peppers under drip irrigation and plastic mulch, 72(4), 575.

*Requisitos de riego para plátano en siete regiones ecológicas de Puerto Rico, 72(4), 599.

*Requisitos de riego para sorgo en las costas sur y norte de Puerto Rico, 72(4), 585.

*Seasonal consumptive use of water by Bell and Cubanelle peppers in semiarid and humid coastal sites in Puerto Rico, 72(4), 609.

Tres niveles de riego y crecimiento y rendimiento del maíz dulce (*Zea mays* cv. Suresweet) regado por goteo, 72(4), 565.

Green, J.
Yam protection. II. Antracnose, yield, and profit of monocultures and interplanting, 72(2), 179.

Gúzman-Pérez, C. D.
Allelopathic effects of seven weed species on pumpkin (*Curcubita moschata*) under greenhouse conditions, 72(3), 491 (RN).

Eggplant (*Solanum melongena* L.) transplants and exudates from roots of seven weeds, 72(3), 495 (RN).

Hepperly, P. R.
Nutritional influences on *Volvariella volvacea* (Bull. ex. Fr.) Sing. growth in Puerto Rico. I. Carbon and Nitrogen, 72(1), 19.

†Sorghum rust, 72(1), 65.

Fungi and insect damage to soybean seeds harvested at immature stages in tropical environments. 72(1), 73.

Yam protection. II. Antracnose, yield, and profit of monocultures and interplanting, 72(2), 179.

Agronomic performance of hard endosperm opaque-2 corn populations in Puerto Rico, 72(2), 309.

†*Coniothyrium* leafspot: A newly described disease of sorghum, 72(2), 319 (RN).

Hernández-López, E.
Susceptibilidad de 53 cultivares de café, *Coffea arabica*, al daño del minador del café, *Leucoptera coffeella*, en Puerto Rico, 72(1), 165 (RN).

Irizarry, H.
*Nutrient uptake and dry matter composition in the plant crop and first ratoon of the Grand Nain banana grown on an Ultisol, 72(3), 337.

*The Lacknau plantain: A high yielding cultivar with field resistance to the corm weevil, *Cosmopolites sordidus* (Germar), 72(3), 353.

Kluson, R. A.
Rhizobium inoculation of *Phaseolus vulgaris* in Lajas, Puerto Rico, 72(3), 427.

Lee, C. T.
†Effect of plant spacing on growth and yield of winged bean, 72(2), 273.

Licha, M.
Tanier (*Xanthosoma* spp.) propagation in vitro, 72(3), 413.

Liu, L. C.
Período crítico de competencia de las malezas en el tomate de trasplante, 72(2), 291.

*Optimum time interval and frequency of glyphosate application for weed control in plantain (*Musa* sp.), 72(2), 297.

*Chemical control of climbing mimosa in a local pasture, 72(4), 615 (RN).

Liu, L. J.
*Tanier (*Xanthosoma* spp.) propagation in vitro, 72(3), 413.

López, J.
Dwarfing effect of interstems and growth and yield components of mango, 72(4), 501.

Luciano, B.
Determinate pigeonpeas [*Cajanus cajan* (L) Mills] genotype on the north and south coasts of Puerto Rico, 72(4), 545.

*Senior author.
†Sole author.
RN Research note.

Lugo, M. L.
*Período crítico de competencia de las malezas en el tomate de trasplante, 72(2), 291.

Lugo-Mercado, H. M.
*Yield response of vegetables to tillage techniques on a Vertisol of the semiarid region of Puerto Rico, 72(1), 119.

Maldonado-Capriles, J.
Contribution to the knowledge of *Rasahus* Amyot and Serville, 1843 (Hemiptera:Reduviidae): description of a new species, 72(2), 247.

Contribution to the knowledge of American Ectrichodiinae. I. Notes about *Rhiginia* (Hemiptera:Reduviidae), 72(2), 251.

Martin, F. W.
*Sensory evaluation of cooked new sweet potato selections, 72(2), 265.

†Planting dates and morphology and growth of four sweet potato clones in Puerto Rico, 72(3), 437.

Mateo-Solano, M.
*Heritabilities and phenotypic correlations for seed yield and seed yield components of beans, 72(2), 301.

Medina-Gaud, S.
Thrips palmi (Thysanoptera:Thripidae): A new insect pest for Puerto Rico, 72(2), 327 (RN).

False smut, a new disease record from Puerto Rico's rice fields, 72(2), 331 (RN).

Lissorhoptrus isthmicus Kuschel (Coleoptera:Curculionidae): A new rice pest for Puerto Rico, 72(2), 333 (RN).

Host plants of pentatomids affecting rice fields in Puerto Rico, 72(3), 365.

Natural enemies of pentatomids affecting rice fields in Puerto Rico, 72(3), 371.

Méndez-Cruz, A. V.
*Yield and nutritive value of hay from five tropical grasses at three harvesting intervals, 72(1), 109.

Production of grazed tropical grasses in different agroecosystems in Puerto Rico. III. Semiarid, 72(2) 211.

*Storage and digestibility, voluntary intake and chemical components of hay of five tropical grasses, 72(4), 531.

Mendoza, N.
Determinate pigeonpeas [*Cajanus cajan* (L) Mills] genotype on the north and south coasts of Puerto Rico, 72(4), 545.

Michelini, S.
*Vesicular-arbuscular mycorrhizal fungus inoculation of sour orange seedlings in Barbados, 72(2), 191.

Mignucci, J. S.
*Yam production. II. Antracnose, yield, and profit of monocultures and interplantings, 72(2), 179.

Mohammed, S.
Some constituents of mabí bark, 72(4), 625 (RN).

Morales, A.
*Determinate pigeon peas [*Cajanus cajan* (L) Mills] genotype on the north and south coasts of Puerto Rico, 72(4), 545.

Murphy, N. F.
Pilot plant fermentation of high test molasses compared with blackstrap molasses using factorial design, 72(1), 9.

*Evaluation of a commercial yeast nutrient for high test molasses fermentation, 72(1), 173 (RN).

The use of distillers' dry yeast as inoculum for rum fermentation in pilot plant scale, 72(3), 445.

†Batch fermentation patterns for different strains of *Zymomonas* in high test molasses and blackstrap molasses, 72(3), 475 (RN).

†Ethanol production from blackstrap molasses by *Zymomonas mobilis* and *Saccharomyces* sp., 72(3), 483 (RN).

†*Zymomonas mobilis* batch and fed-batch fermentation of high test molasses, 72(3), 485 (RN).

Pilot plant scale evaluation of a commercial yeast nutrient for high test molasses fermentation, 72(4), 509.

Nemec, S.
Vesicular-arbuscular mycorrhizal fungus inoculation of sour orange seedlings in Barbados, 72(2), 191.

Oramas, D.
The Lacknau plantain: A high yielding cultivar with field resistance to the corm weevil, *Cosmopolites sordidus* (Germar), 72(3), 353.

Orengo, E.
Yield response of vegetables to tillage techniques on a Vertisol of the semiarid region of Puerto Rico, 72(1), 119.

*Senior author.
†Sole author.
RN Research note.

Ortiz, A. A.
Bacteria associated with the West Indian fruit fly *Anastrepha obliqua* (Macquart) (Diptera:Tephritidae), 72(3), 489 (RN).

Ortiz, C.
*Fungi and insect damage to soybean seeds harvested at immature stages in tropical environments, 72(1), 73.

Yield response of vegetables to tillage techniques on a Vertisol of the semiarid region of Puerto Rico, 72(1), 119.

Ortiz, F. H.
Determinate pigeonpeas [*Cajanus cajan* (L) Mills] genotype on the north and south coasts of Puerto Rico, 72(4), 545.

Pantoja, A.
**Thrips palmi* (Thysanoptera:Thripidae): A new insect pest for Puerto Rico, 72(2), 327 (RN).

*False smut, a new disease record from Puerto Rico's rice fields, 72(2), 331 (RN).

**Lissorhoptrus isthmicus* Kuschel (Coleoptera:Curculionidae): A new rice pest for Puerto Rico, 72(2), 333 (RN).

Host plants of pentatomids affecting rice fields in Puerto Rico, 72(3), 365.

Natural enemies of pentatomids affecting rice fields in Puerto Rico, 72(3), 371.

Sequential sampling plan, yield-loss components and economic thresholds for the pepper weevil, *Anthonomus eugenii* Cano (Coleoptera:Curculionidae), 72(3), 375.

Evaluation of relative sampling methods for population estimation of the pepper weevil, *Anthonomus eugenii* Cano (Coleoptera: Curculionidae), 72(3), 387.

Pérez, A.
Effect of dwarfing rootstocks on tree size and yield of selected mango varieties, 72(1), 1.

*Dwarfing effect of interstems and growth and yield components of mango, 72(4), 501.

Persaud, T.
*Cost of a drip irrigation system for vegetable farming in Puerto Rico, 72(1), 31.

Labor-input requirements for experimental production of drip irrigated vegetables, 72(1), 41

Quiles-Belén, A.
*Corn response to N applications and population densities at two locations in Puerto Rico, 72(1), 127.

Ramírez-Oliveras, G.
Yield and nutritive value of hay from five tropical grasses at three harvesting intervals, 72(1), 109.

Randel, P. F.
*Confinement feeding of dairy cows based on stargrass as green chopped fodder or hay, 72(2), 231.

Recio de Hernández, E.
Freezing and shelf life of sweet potato and citron bars, 72(1), 171 (RN).

Reyes-Soto, I.
Effect of dwarfing rootstocks on tree size and yield of selected mango varieties, 72(1), 1.

Dwarfing effect of interstems on growth and yield components of mango, 72(4), 501.

Rivera, E.
Nutrient uptake and dry matter composition in the plant crop and first ratoon of the Grand Nain banana grown on an Ultisol, 72(3), 337.

The Lacknau plantain: A high yielding cultivar with field resistance to the corm weevil, *Cosmopolites sordidus* (Germar), 72(3), 353.

Rivera, L. E.
Labor-input requirements for experimental production of drip irrigated vegetables, 72(1), 41.

Root distribution of nitrogen fertigated sweet peppers under drip irrigation, 72(1), 51.

Dry beans and varying cultural practices at two locations on the coastal plain of Puerto Rico, 72(4), 521.

Nutrient uptake and growth characteristics of nitrogen fertigated sweet peppers under drip irrigation and plastic mulch, 72(4), 575.

Rodríguez, J.
Nutrient uptake and dry matter composition in the plant crop and first ratoon of the Grand Nain banana grown on an Ultisol, 72(3), 337.

Rodríguez, J. A.
The Lacknau plantain: A high yielding cultivar with field resistance to the corm

*Senior author.
RN Research note.

weevil, *Cosmopolites sordidus* (Germar), 72(3), 353.

Rodríguez T.
Shelf life and acceptability of hot water-treated mangoes, 72(3), 469.

Rodríguez de Cianzio, S.
Fungi and insect damage to soybean seeds harvested at immature stages in tropical environments, 72(1), 73.

Rodríguez-García, J.
Optimum time interval and frequency of glyphosate application for weed control in plantain (*Musa* sp.), 72(2), 297.

Román-Hernández, O.
Dry beans and varying cultural practices at two locations on the coastal plain of Puerto Rico, 72(4), 521.

Rosa-Márquez, E.
Tanier (*Xanthosoma* spp.) propagation in vitro, 72(3), 413.

Ruiz, H.
Thrips palmi (Thysanoptera:Thripidae): A new insect pest for Puerto Rico, 72(2), 327 (RN).

Saladín-García, F.
Heritabilities and phenotypic correlations for seed yield and seed yield components of beans, 72(2), 301.

Schroder, E. C.
Rhizobium inoculation of *Phaseolus vulgaris* in Lajas, Puerto Rico, 72(3), 427.

Seaforth, C. E.
*Some constituents of mabí bark, 72(4), 625 (RN).

Segarra, A.
Chemical control of *Liriomyza sativae* and of Lepidoptera larvae on tomato, 72(1), 169 (RN).

Thrips palmi (Thysanoptera:Thripidae): A new insect pest for Puerto Rico, 72(2), 327 (RN).

Segarra, A. E.
†Identity and economic importance of fruit flies attacking mango in Puerto Rico: A two-year survey, 72(2), 325 (RN).

*Sequential sampling plan, yield-loss components and economic thresholds for the pepper weevil, *Anthonomus eugenii* Cano (Coleoptera:Curculionidae), 72(3), 375.

*Evaluation of relative sampling methods for population estimation of the pepper weevil, *Anthonomus eugenii* Cano (Coleoptera:Curculionidae), 72(3), 387.

Segarra-Carmona, A. E.
*Relative susceptibility of *Crotalaria* spp. to attack by *Etiella zinckenella* in Puerto Rico, 72(1), 147.

*Notes on the natural enemies of *Etiella zinckenella* in Puerto Rico, 72(1), 153.

*The sorghum midge, *Contarinia sorghicola*, a sorghum pest newly introduced to Puerto Rico, 72(1), 161 (RN).

*Bacteria associated with the West Indian fruit fly *Anastrepha obliqua* (Macquart) (Diptera:Tephritidae), 72(3), 489 (RN).

Semidey-Laracuente, N.
Evaluation of diquat for postemergence weed control in pumpkin (*Curcubita moschata*), 72(2), 285.

Allelopathic effects of seven weed species to pumpkin (*Curcubita moschata*) under greenhouse conditions, 72(3), 491 (RN).

Eggplant (*Solanum melongena* L.) transplants and exudades from roots of seven weeds, 72(3), 495 (RN).

Siberio-Torres, V.
Yield and nutritive value of hay from five tropical grasses at three harvesting intervals, 72(1), 109.

Storage and digestibility, voluntary intake and chemical components of hay of five tropical grasses, 72(4), 531.

Sievens, G.
Chemical control of climbing mimosa in a local pasture, 72(4), 615 (RN).

Silva-Negrón, L.
Refrigeration before incubation and its effect on recovery of two types of bacteria from food, 72(4), 617 (RN).

Sotomayor-Ríos, A.
Corn response to N applications and population densities at two locations in Puerto Rico, 72(1), 127.

Tergas, L. E.
*Response of tropical grasses to potassium fertilization in the humid coastal region of Puerto Rico, 72(1), 81.

*Production of grazed tropical grasses at Toa Baja in the humid northern coastal plains of Puerto Rico, 72(1), 91.

*Production of grazed tropical grasses in different agroecosystems in Puerto Rico. I. Humid mountain, 72(1), 99.

*Senior author.
†Sole author.
RN Research note.

*Production of grazed tropical grasses in different agroecosystems in Puerto Rico. II. Humid northern coastal plains, 72(2), 201.

*Production of grazed tropical grasses in different agroecosystems in Puerto Rico. III. Semiarid, 72(2), 211.

*Forage production of nine alfalfa cultivars in the semiarid region of Puerto Rico, 72(2), 221.

Torres-Cardona, S.
Corn response to N applications and population densities at two locations in Puerto Rico, 72(1), 127.

Torres-López, R.
Yam production. II. Antracnose, yield, and profit of monocultures and interplanting, 72(2), 179.

Torres-López, R. I.
*Nutritional influences on *Volvariella volvacea* (Bull. ex. Fr.) Sing. growth in Puerto Rico. I. Carbon and nitrogen, 72(1), 19.

Torres-Rivera, C. N.
Bacteria associated with the West Indian fruit fly *Anastrepha obliqua* (Macquart) (Diptera:Tephritidae), 72(3), 489 (RN).

Unander, D. W.
*Pepper cultivars on the northwestern coast of Puerto Rico, 72(3), 395.

Varela-Ramírez, F.
Pepper cultivars on the northwestern coast of Puerto Rico, 72(3), 395.

Velázquez, Y. A.
**Rhizobium* inoculation of *Phaseolus vulgaris* in Lajas, Puerto Rico, 72(3), 427.

Vélez-Colón, R.
*Evaluación de variedades de uva de mesa (*Vitis vinifera* L.) cosechadas en la subestación de Fortuna en 1986, 72(2), 335 (RN).

Vélez-Santiago, J.
Response of tropical grasses to potassium fertilization in the humid coastal region of Puerto Rico, 72(1), 81.

Production of grazed tropical grasses at Toa Baja in the humid northern coastal plains of Puerto Rico, 72(1), 91.

Production of grazed tropical grasses in different agroecosystems in Puerto Rico. I. Humid mountain, 72(1), 99.

Production of grazed tropical grasses in different agroecosystems in Puerto Rico. II. Humid northern coastal plains, 72(2), 201.

Production of grazed tropical grasses in different agroecosystems in Puerto Rico. III. Semiarid, 72(2), 211.

Forage production of nine alfalfa cultivars in the semiarid region of Puerto Rico, 72(2), 221.

Vera de Saldaña, D.
Response of tropical grasses to potassium fertilization in the humid coastal region of Puerto Rico, 72(1), 81.

Production of grazed tropical grasses at Toa Baja in the humid northern coastal plains of Puerto Rico, 72(1), 91.

Production of grazed tropical grasses in different agroecosystems in Puerto Rico. I. Humid mountain, 72(1), 99.

Production of grazed tropical grasses in different agroecosystems in Puerto Rico. II. Humid northern coastal plains, 72(2), 201.

Forage production of nine alfalfa cultivars in the semiarid region of Puerto Rico, 72(2), 221.

Wessel-Beaver, L.
*Agronomic performance of hard endosperm opaque-2 corn populations in Puerto Rico, 72(2), 309.

*Senior author.
RN Research note.

SUBJECTS

Achrysonocharoides sp.
Coffee leaf miner parasitoid in Puerto Rico, 72(2), 255.
Alfalfa, 9 cultivars
Forage production in semiarid region, 72(2), 221.
Allelopathic effects of weeds
Pumpkin, 72(3), 491.
Amaranthus dubius
Allelopathic effect on pumpkin, 72(3), 491.
Exudates from its roots in eggplants, 72(3), 495.
Anastrepha obliqua
Bacteria associated, 72(3), 489.
Anastrepha sp.
Identity and economic importance, 72(2), 325.
Anthonomus eugenii
Sampling methods for population estimation, 72(3), 387.
Yield-loss components and economic thresholds, 72(3), 375.
Anthracnose
Damage to immature soybean seeds, 72(1), 73.
Yield and profit in monocultures and interplantings of yams, 72(2), 179.
Asparagus
Seed production, 72(4), 619.
Bacillus coagulans
Recovery from food refrigerated before incubation, 72(4), 617.
Bacillus maurans
Recovery from food refrigerated before incubation, 72(4), 617.
Bacteria
Recovery from food refrigerated before incubation, 72(4), 617.
Banana
Dry matter composition, 72(3), 337.
Nutrient uptake, 72(3), 337.
Bean
Yield response to tillage techniques, 72(1), 119.
Bean, winged
Plant spacing and growth and yield, 72(2), 273.
Beans
Inoculation with *Rhizobium*, 72(3), 427.
Beans, dry
Varying cultural practices at two locations, 72(4), 515.
Beans, seed yields
Heritabilities and phenotypic correlations, 72(2), 301.
Beskia aelops
Natural enemy of pentatomids, 72(3), 371.
Brachiaria brizantha
Production in the humid northern coastal plains, 72(1), 91.
Brachiaria humidicola
Production in the humid mountains, 72(1), 99.
Brachiaria ruziziensis
Production in the humid northern coastal plains, 72(1), 91.
Café
Susceptibilidad de cultivares al minador, 72(1), 165.
Cajanus cajan
Determinate genotypes, 72(4), 545.
Light trap studies of *Heliothis virescens* and *Etiella zinckenella*, 72(4), 557.
Some host plants of its leafhopper, 72(3), 499.
Callie, hay
Storage and digestibility, voluntary intake and chemical components, 72(4), 531.
Capsicum annuum
Yield on northwestern coast, 72(3), 395.
Cercospora kikuchii
Damage to immature soybean seeds, 72(1), 73.
Chrysonotomya sp.
Coffee leaf miner parasitoid in Puerto Rico, 72(2), 255.
Cirrospiloides sp.
Coffee leaf miner parasitoid in Puerto Rico, 72(2), 255.
Citron, bars
Freezing and shelf life, 72(1), 171.
Citrus medica, bars
Freezing and shelf life, 72(1), 171.
Cocoyam, in vitro cultured tissue
Salt and sugar levels for effective plant regeneration, 72(4), 515.

Coffee
Bacterial leaf spot, 72(4), 621.
Susceptibility of cultivars to leaf miner, 72(1), 165.
Coffea arabica
Distribution of leaf miner and its parasitoids in tree canopy, 72(1), 141.
Coffee leaf miner
Distribution in the tree canopy, 72(1), 141.
Parasitoids in Puerto Rico, 72(2), 255.
Coleoptera
A new rice pest for Puerto Rico, 72(2), 333.
Colletotrichum dematium
Damage to immature soybean seeds, 72(1), 73.
Colletotrichum gloeosporioides
Yield and profit in monocultures and interplantings of yams, 72(2), 179.
Coniothyrium leaf spot
A newly described disease of sorghum, 72(2), 319.
Consumptive use, seasonal
Bell and Cubanelle peppers, 72(4), 609.
Contarinia sorghicola
A newly introduced pest to Puerto Rico, 72(1), 161.
Corn
Yield response to tillage techniques, 72(1), 119.
Corn, Opaque-2 populations
Agronomic performance, 72(2), 309.
Corn, sweet
Estimation of irrigation requirements, 72(2), 277.
Corn, two locations
N applications and population densities, 72(1), 127.
Cosmopolites sordidus
Resistance of Lacknau plantain cultivar, 72(3), 353.
Cows, dairy, confinement feeding
Green chopped stargrass or hay, 72(2), 231.
Crotalaria
Susceptibility to *Etiella zinckenella*, 72(1), 147.
Crotalaria palida
Hymenopterous parasitoid of larvae feeding on, 72(1), 153.
Curcubita moschata
Allelopathic effects of weeds, 72(3), 491.
Diquat as a postemergence herbicide, 72(2), 285.
Curculionidae
A new rice pest for Puerto Rico, 72(2), 333.
Cynodon dactylon
Production in humid northern coastal plains (Barceloneta), 72(2), 201.
Production in semiarid region, 72(2), 211.
Production in the humid mountains, 72(1), 99.
Production in the humid northern coastal plains (Toa Baja), 72(1), 91.
Response to potassium fertilization, 72(1), 81.
Cynodon dactylon, hay
Storage and digestibility, voluntary intake and chemical components, 72(4), 531.
Yield and nutritive value at three harvesting intervals, 72(1), 109.
Cynodon nlemfuensis
Confinement feeding of dairy cows, 72(2), 231.
Production in humid northern coastal plains (Barceloneta), 72(2), 201.
Production in semiarid region, 72(2), 211.
Production in the humid mountains, 72(1), 99.
Production in the humid northern coastal plains (Toa Baja), 72(1), 91.
Response to potassium fertilization, 72(1), 81.
Cynodon nlemfuensis, hay
Yield and nutritive value at three harvesting intervals, 72(1), 109.
Cynodon plectostachyus
Production in humid northern coastal plains (Barceloneta), 72(2), 201.
Production in semiarid region, 72(2), 211.
Production in the humid mountains, 72(1), 99.
Production in the humid northern coastal plains (Toa Baja), 72(1), 91.
Storage and digestibility, voluntary intake and chemical components, 72(4), 531.
Cynodon plectostachyus, hay from
Yield and nutritive value at three harvesting intervals, 72(1), 109.
Cynodon sp.
Production in humid northern coastal plains, 72(2), 201.
Dichanthium annulatus
Production in semiarid region, 72(2), 211.
Digitaria decumbens
Production in the humid mountains, 72(1), 99.
Production in the humid northern coastal plains (Barceloneta), 72(2), 201.
Production in the humid northern coastal plains (Toa Baja), 72(1), 91.

Response to potassium fertilization, 72(1), 81.
Digitaria milanjiana
Production in humid northern coastal plains, 72(2), 201.
Digitaria pentzii
Production in humid northern coastal plains, 72(2), 201.
Production in the humid mountains, 72(1), 99.
Response to potassium fertilization, 72(1), 81.
Digitaria pentzii, hay
Storage and digestibility, voluntary intake and chemical components, 72(4), 531.
Yield and nutritive value at three harvesting intervals, 72(1), 109.
Dioscorea spp.
Anthracnose, yield and profit, 72(2), 179.
Diquat
Post emergence herbicide in pumpkin, 72(2), 285.
Dwarfing interstems in trees
Growth and yield components in mango, 72(4), 501.
Echinochloa colonum
Allelopathic effect on pumpkin, 72(3), 491.
Exudates from its roots in eggplants, 72(3), 495.
Ectrichodiinae
Notes about *Rhiginia* sp, 72(2), 251.
Eggplant
Yield response to tillage techniques, 72(1), 119.
Eggplant transplants
Exudates of roots of weeds, 72(3), 495.
Elevation
Relation to temperature, 72(3), 449.
Empoasca spp.
Some host plants, 72(3), 499.
Endosperm, hard
Performance of Opaque-2 populations in Puerto Rico, 72(2), 309.
Eriochloa polystachya
Production in the humid northern coastal plains, 72(1), 91.
Ethanol
Production from blackstrap molasses by *Zymomonas mobilis* and *Saccharomyces*, 72(3), 483.
Etiella zinckenella
Damage to immature soybean seeds, 72(1), 73.
Natural enemies, 72(1), 153.
Susceptibility of *Crotalaria* spp. to its attack, 72(1), 147.
Etiella zinckenella on pigeon pea
Light trap studies, 72(4), 557.
Euphorbia heterophylla
Allelopathic effect on pumpkin, 72(3), 491.
Exudates from its roots in eggplants, 72(3), 495.
Evapotranspiration, potential
Hargreaves-Samani technique, 72(1), 57.
Vieques Island, 72(1), 177.
Fascioliasis
Relationship to watercress cultivation, 72(3),405.
Fermentation batch
Patterns for strains of *Zymomonas*, 72(3), 475.
Fermentation, batch and fed-batch
High test molasses by *Zymomonas mobilis*, 72(3), 485.
Fermentation, high test molasses
Commercial yeast nutrient, 72(1), 173.
Commercial yeast nutrient evaluation, 72(4), 509.
Compared with blackstrap molasses, 72(1), 9.
Flies, fruit
Identity and economic importance, 72(2), 325.
Fly, fruit, West Indian
Bacteria associated, 72(3), 489.
Fodder, hay, stargrass
Confinement feeding of dairy cows, 72(2), 231.
Forage, alfalfa
Production of 9 cultivars in semiarid region, 72(2), 221.
Fruit fly, West Indian
Bacteria associated, 72(3), 489.
Fungi, vesicular-arbuscular
Inoculation of sour orange seedlings, 72(2), 191.
Glomus spp.
Inoculation of sour orange seedlings, 72(2), 191.
Glyphosate
Optimum time interval and frequency of application, 72(2), 297.
Grapes, table
Variety evaluation at Fortuna, 72(2), 335.
Grasses, tropical
Response to potassium fertilization, 72(1), 81.
Grasses, tropical, grazed
Production in humid mountains, 72(1), 99.
Production in humid northern coastal plains (Barceloneta), 72(2), 201.
Production in humid northern coastal plains (Toa Baja), 72(1), 91.

Production in semiarid region, 72(2), 211.
Grasses, tropical, hay
Storage and digestibility, voluntary intake and chemical components, 72(4), 531.
Grasses, tropical, hay from
Yield and nutritive value at three harvesting intervals, 72(1), 109.
Guinea grass, hay
Storage and digestibility, voluntary intake and chemical components, 72(4), 531.
Gymnoclytia
Natural properties of pentatomids, 72(3), 371.
Hargreaves and Samani
Model for potential evapotranspiration estimation, 72(1), 177.
Hargreaves-Samani technique
Potential evapotranspiration for the south coast, 72(1), 57.
Hay, fodder, stargrass
Confinement feeding of dairy cows, 72(2), 231.
Hay, tropical grasses
Storage and digestibility, voluntary intake and chemical components, 72(4), 531.
Yield and nutritive value at three harvesting intervals, 72(1), 109.
Heliothis virescens on pigeonpea
Light trap studies, 72(4), 557.
Hemiptera
Description of new species of *Rasahus*, 72(2), 247.
Description of *Rhiginia dureti* Carpintero from Peru, 72(2), 251.
Heritabilities and phenotypic correlations
Bean seed yield and yield components, 72(2), 301.
Heterospilus etiellae
Larval parasitoid, 72(1), 153.
Horismenus sp.
Coffee leaf miner parasitoid in Puerto Rico, 72(2), 255.
Iconella etiellae, larval parasitoid
Recovered from soybean, 72(1), 153.
Inoculation, sour orange seedlings
Vesicular-arbuscular mycorrhizal fungi, 72(2), 191.
Inoculum, distillers' dry yeast
Rum fermentation on pilot plant scale, 72(3), 445.
Interstems in mango
Dwarfing effect on growth and yield, 72(4), 501.
Ipomoea batatas
Planting dates, morphology and growth, 72(3), 437.
Sensory evaluation, 72(2), 265.
Ipomoea batatas, bars
Freezing and shelf life, 72(1), 171.
Irrigation, drip
Cost for vegetable farming, 72(1), 31.
Labor input in vegetable production, 72(1), 41.
Root distribution of fertigated peppers, 72(1), 51.
Irrigation (drip) and plastic mulch
Nutrient uptake in sweet peppers, 72(4), 575.
Irrigation of sweet corn
Estimation requirements, 72(2), 277.
Leaf miner, coffee
Distribution of parasitoids in the tree canopy, 72(1), 141.
Susceptibility of coffee cultivars, 72(1),165.
Leaf spot
A *Coniothyrium* disease in sorghum, 72(2), 319.
Leaf spot, bacteria
Coffee, 72(4), 621.
Leafhopper, pigeon pea
Some host plants, 72(3), 499.
Lepidoptera, larvae
Chemical control in tomato, 72(1), 169.
Leucoptera coffeella
Distribution of parasitoids in the tree canopy, 72(1), 141.
Parasitoids in Puerto Rico, 72(2), 255.
Susceptibility of coffee cultivars, 72(1), 165.
Lima bean pod borer
Susceptibility of *Crotalaria* spp to its attack, 72(1), 147.
Liriomyza sativae
Chemical control in tomato, 72(1) , 169.
Lissorhoptrus isthmicus
A new rice pest for Puerto Rico, 72(2), 333.
Lycopersicon esculentum
Chemical control of *Liriomyza sativae* and Lepidoptera larvae, 72(1), 169.
Mabí, bark
Chemical constituents, 72(4), 625.
Maíz dulce
Irrigation requirements, estimation of, 72(2), 277.
Requisitos de riego, estimación de, 72(2), 277.
Tres niveles de riego, crecimiento y rendimiento, 72(4), 565.

Mango
Fruit flies which attack, 72(2), 325.
Mango, dwarfing rootstocks
Tree size and yield of varieties, 72(1), 1.
Mango, growth and yield
Dwarfing effect of interstems, 72(4), 501.
Mangoes, hot water treated
Shelf life and acceptability, 72(3), 469.
Mangoes, 3 cultivars
Shelf life when hot water-treated, 72(3), 469.
Medicago sativa, 9 cultivars
Forage production in semiarid region, 72(2), 221.
Mimosa, climbing
Chemical control in a local pasture, 72(4), 615.
Mirax insularis
Coffee leaf miner parasitoid in Puerto Rico, 72(2), 255.
Parasitoid of coffee leaf miner, 72(1), 141.
Molasses, blackstrap
Ethanol production by *Zymomonas mobilis* and *Saccharomyces* sp, 72(3), 483.
Molasses high test
Batch and fed-batch fermentation by *Zymomonas mobilis*, 72(3), 485.
Commercial yeast nutrient for fermentation, 72(1), 173.
Pilot plant scale evaluation of a commercial yeast nutrient in fermentation, 72(4), 509.
Molasses, high test, fermentation
Compared with blackstrap, 72(1), 9.
Molasses, high test vs blackstrap
Batch fermentation patterns for strain of *Zymomonas*, 72(3), 475.
Mormidea angustata Stål
Host plants in rice fields, 72(3), 365.
Pentatomid parasitized by natural enemies, 72(3), 371.
Musa sp.
Optimum time interval and frequency of glyphosate application, 72(2), 297.
Mushroom, straw
Carbon and nitrogen nutritional influences on growth, 72(1), 19.
Nasturtium officinale
Relationship of cultivation sites to fascioliasis incidence, 72(3), 405.
Nezara viridula
Damage to immature soybean seeds, 72(1), 73.
Nitrogen applications
Corn at two locations, 72(1), 127.
Oebalus ypsilon-griseus
Host plants in rice fields, 72(3), 365.
Pentatomid parasitized by natural enemies, 72(3), 371.
Ooencyrtus submetallicus
Natural enemy of pentatomids, 72(3), 371.
Opaque-2 corn populations
Agronomic performance, 72(2), 309.
Orange, sour, seedlings
Inoculation with vesicular-arbuscular mycorrhizal fungi, 72(2), 191.
Panicum maximum
Production in humid northern coastal plains (Barceloneta), 72(2), 201.
Production in semiarid region, 72(2), 211.
Production in the humid mountains, 72(1), 99.
Production in the humid northern coastal plains (Toa Baja), 72(1), 91.
Storage and digestibility, voluntary intake and chemical components, 72(4), 531.
Panicum maximum, hay
Yield and nutritive value at three harvesting intervals, 72(1), 109.
Paraquat
Postemergence weed control in pumpkin, 72(2), 285.
Parthenium hysterophorus
Allelopathic effect on pumpkin, 72(3), 491.
Exudates from its roots in eggplants, 72(3), 495.
Pentatomids
Host plants in rice fields, 72(3), 365.
Natural enemies in rice fields, 72(3), 371.
Pepper
Yield response to tillage technique, 72(1), 119.
Pepper cultivars
Yield on northwestern coast, 72(3), 395.
Pepper weevil
Sampling methods for population estimation, 72(3), 387.
Yield-loss components and economic thresholds, 72(3), 375.
Peppers
Seasonal consumptive use of water, 72(4), 609.
Peppers, sweet
Root distribution of nitrogen fertigated, 72(1), 51.
Peppers, sweet, nitrogen fertigated
Nutrient uptake and growth characteristics, 72(4), 575.

Peronospora manshurica
Damage to immature soybean seeds, 72(1), 73.
Phaseolus vulgaris
Heritabilities and phenotypic correlations, 72(2), 301.
Inoculation with *Rhizobium*, 72(3), 427.
Varying cultural practices at two locations, 72(4), 521.
Phenotypic correlations
Bean seed yield and yield components, 72(2), 301.
Phomopsis spp.
Damage to immature soybean seeds, 72(1), 73.
Pigeon pea
Determinate genotypes, 72(4), 545.
Light trap studies of *Heliothis virescens* and *Etiella zinckenella*, 72(4), 557.
Some host plants of its leafhopper, 72(3), 499.
Plantain
Irrigation requirements, 72(4), 599.
Optimum time interval and fequency of glyphosate application, 72(2), 297.
Plantain, Lacknau cultivar
Resistance to corm weevil, 72(3), 353.
Plátano
Requisitos de riego, 72(4), 599.
Potassium fertilization
Response of tropical grasses in the humid coastal region, 72(1), 81.
Pseudomonas syringae
Leaf spot in coffee, 72(4), 621.
Psophocarpus tetragonolobus
Plant spacing, growth, yield, 72(2), 273.
Puccinia purpurea
Sorghum, 72(1), 65.
Pumpkin
Allelopathic effects of weeds, 72(3), 491.
Diquat as a postemergence herbicide, 72(2), 285.
Rasahus sp.
Description of new species, 72(2), 247.
Reduviidae
Description of new species of *Rasahus*, 72(2), 247.
Description of *Rhiginia dureti* Carpintero from Peru, 72(2), 251.
Rhiginia sp.
New species described from Peru, 72(2), 251.
Rhizobium
Inoculation of *Phaseolus vulgaris*, 72(3), 427.
Rice
Attacked by *Lissorhoptrus isthmicus* in Puerto Rico, 72(2), 333.
New disease record from Puerto Rico, 72(2), 331 (RN).
Rice fields
Host plants of pentatomids, 72(3), 365.
Natural enemies of pentatomids, 72(3), 371.
Riego de maíz dulce
Estimación de los requisitos, 72(2), 277.
Root distribution, sweet peppers
Nitrogen fertigated and drip irrigated, 72(1), 51.
Rottboelia exaltata
Allelopathic effect on pumpkin, 72(3), 491.
Exudates from its roots in eggplants, 72(3), 495.
Rootstocks, dwarfing
Tree size and yield of varieties, 72(1), 1.
Rum fermentation
Distillers' dry yeast inoculum, 72(3), 445.
Rust, sorghum, 72(1), 65.
Saccharomyces
Ethanol production from blackstrap molasses, 72(3), 483.
Santainezia bicolor
Synonym of *Rhiginia haenschi*, 72(2), 251.
Seed asparagus
Production in Puerto Rico, 72(4), 619.
Seed, yield and yield components
Heritabilities and phenotypic correlations in beans, 72(2), 301.
Seedlings, orange, sour
Inoculation with vesicular-arbuscular mycorrhizal fungi, 72(2), 191.
Shelf-life and acceptability
Hot water-treated mangoes, 72(3), 469.
Smut, false
A new disease record from Puerto Rico's rice fields, 72(2), 331.
Solanum melongena
Exudates of roots of weeds, 72(3), 495.
Sorghum
A newly described disease caused by *Coniothyrium*, 72(2), 319.
Irrigation requirements in the south and north coasts, 72(4), 585.
Sorghum halepense
Allelopathic effect on pumpkin, 72(3), 491.
Exudates from its roots in eggplants, 72(3), 495.
Sorghum midge
A newly introduced pest to Puerto Rico, 72(1), 161.
Sorghum rust, 72(1), 65.

Sorgo
Requisitos de riego en las costas sur y norte, 72(4), 585.
Soybean seeds, immature
Fungi and insect damage, 72(1), 73.
Spacing
Growth and yield of winged bean, 72(2), 273.
Stargrass, green chopped fodder or hay
Confinement feeding of dairy cow, 72(2), 231.
Sweet potato, bars
Freezing and shelf life, 72(1), 171.
Sweet potato, 4 clones
Planting dates, morphology and growth, 72(3), 437.
Sweet potato selections, cooked
Sensory evaluation, 72(2), 265.
Tanier
In vitro propagation, 72(3), 413.
Temperature
Relation to elevation, 72(3), 449.
Thrips palmi
A new insect pest in Puerto Rico, 72(2), 327.
Tillage techniques
Yield response of vegetables in semiarid region, 72(1), 119.
Tissue culture, cocoyam
Salt and sugar levels for effective plant regeneration, 72(4), 515.
Tomate
Período crítico de competencia de malezas, 72(2), 291.
Tomato
Chemical control of *Liriomyza sativae* and Lepidoptera larvae, 72(1), 169.
Critical period of weed competition, 72(2), 291.
Yield response to tillage technique, 72(1), 119.
Trianthema portulacastrum
Allelopathic effect on pumpkin, 72(3), 491.
Exudates from its roots in eggplants, 72(3), 495.
Trichogramma sp
Parasitism on *Etiella zinckenella* eggs, 72(1), 153.
Ustilaginoidea oryzae
A new disease record from Puerto Rico's rice fields, 72(2), 331.
Uva de mesa
Evaluación de variedades, 72(2), 335.
Vegetable farming
Cost of drip irrigation system, 72(1), 31.
Vegetables
Yield response to tillage technique, 72(1), 119.
Vegetables, drip irrigated
Labor input requirements, 72(1), 41.
Vieques Island
Potential evapotranspiration with Hargreaves and Samani model, 72(1), 177.
Vitis vinifera
Evaluación de variedades, 72(2), 335.
Variety evaluation at Fortuna, 72(2), 335.
Volvariella volvacea
Carbon and nitrogen nutritional influences on growth, 72(1), 19.
Watercress
Relationship of cultivation sites to fascioliasis incidence, 72(3), 405.
Watermelon
Yield response to tillage technique, 72(1), 119.
Weeds
Critical competition period in tomatos, 72(2), 291.
Exudates from weed roots and eggplant transplants, 72(3), 495.
Weeds, allelopathic effects
Pumpkin, 72(3), 491.
Weevil, corm, plantain
Resistance of Lacknau plantain cultivar, 72(3), 353.
Weevil, pepper
Sampling methods for population estimation, 72(3), 387.
Yield-loss components and economic thresholds, 72(3), 375.
Xanthosoma spp.
In vitro propagation, 72(3), 413.
Yam, monocultures and interplantings Anthracnose, yield and profit, 72(2), 179.
Yeast, commercial
Nutrient for high test molasses fermentation, 72(1), 173.
Yeast, distillers' dry
Inoculum for rum fermentation, 72(3), 445.
Yeast nutrient, commercial
Pilot plant scale evaluation for high test molasses fermentation, 72(4), 509.
Zagrammosoma sp.
Coffee leaf miner parasitoid in Puerto Rico, 72(2), 255.
Zea mays
Estimation of irrigation requirements, 72(2), 277.
N applications and population densities, 72(1), 127.

Zea mays, cv Suresweet
Tres niveles de riego y crecimiento y rendimiento, 72(4), 565.
Zymomonas
Fermentation patterns in high test molasses and blackstrap molasses, 72(3), 475.
Zymomonas mobilis
Batch and fed-batch fermentation of high test molasses, 72(3), 485.
Ethanol production from blackstrap molasses, 72(3), 483.